

Do general relativistic effects limit experiments to test the universality of free fall and the weak equivalence principle?

Anna M. Nobili^{1,2}

¹*Dept. of Physics “E. Fermi”, University of Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy*

²*INFN-Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy*

(Dated: October 26, 2016)

The Universality of Free Fall and the Weak Equivalence Principle, which are at the basis of General Relativity, have been confirmed to 1 part in 10^{13} . Space experiments with macroscopic test masses of different composition orbiting the Earth inside a low altitude satellite aim at improving this precision by two orders of magnitude (with the Microscope satellite, launched on 25 April 2016) and up to four orders of magnitude (with the “Galileo Galilei” - GG satellite). At such a high precision many tiny effects must be taken into account in order to be ruled out as the source of a spurious violation signal. In this work we investigate the general relativistic effects, including those which involve the rotation of both the Earth and the test masses, and show that they are by far too small to be considered even in the most challenging experiment.^a

I. INTRODUCTION

The Universality of Free Fall (UFF), also known as the Weak Equivalence Principle (WEP) refers to the fact that in a gravitational field all bodies fall with the same acceleration regardless of their mass and composition (see e.g. [1]). As stated by Einstein in 1916 [2] the General theory of Relativity (GR) requires this fact to hold.

The best experimental tests so far involve artificial proof masses suspended on rotating torsion balances [3–5] or celestial bodies (Earth and Moon, through Lunar Laser Ranging [6–8]). They show no anomalous differential acceleration between the falling bodies –in the fields of the Earth and the Sun– to about 10^{-13} .

Considerable improvements are expected in the field of the Earth by flying macroscopic proof masses inside a low altitude spacecraft. Microscope [9], launched in April 2016, aims at 10^{-15} , and “Galileo Galilei” (GG) [10] aims at 10^{-17} . As for cold atom tests, at present they have reached a few 10^{-8} [11] and are not expected to compete with high-precision space tests based on macroscopic bodies [12].

Despite their high precision goals, and consequent very small effects to be measured, the GG and Microscope experiments are designed and investigated within Newtonian mechanics, in the assumption that general relativistic effects are too small to compete with the sought for violation signal.

During a recent competitive selection of proposals shortlisted for the medium size mission M4 of the European Space Agency (ESA) this assumption has been questioned by SARP, the “Science Assessment Review Panel” appointed by ESA to evaluate the GG proposal. From their Report on GG we read [13]:

“The breakdown of the WEP is sought in the frame work of the response of test matter to terrestrial Newtonian gravitation. The

source of terrestrial Newtonian gravitation is independent of the Earth’s (non uniform) rotation. Furthermore, the test cylinders in the proposed experiment are spinning. In General Relativity the gravitational field of a spinning source depends on its spin. Also the mass centroid motion of extended spinning test matter in an external gravitational field may depend on its spin and still be geodesic (independent of inertial mass) when its spin is zero. The estimates, based on General Relativity, of the effect of the Earth’s rotation on the motion of each spinning cylinder or the laser interferometer and their relevance to the interpretation of any non null signal at the expected level of accuracy have not been sufficiently explained to the satisfaction of the SARP.”

Experimental evidence of a violation of UFF-WEP would require either a modification/augmentation of GR or the existence of a new composition dependent force of nature. Either way, it would make a revolution in physics. On the other hand, a null result to a very high precision will be a landmark for any attempt at overcoming the current physics impasse. The situation is reminiscent of that at the end of the 19th century, when Michelson and Morley tested by very precise light interferometry the propagation of the newly discovered electromagnetic waves through the ether and proved that the ether does not exist[14]; a very precise null result which in 1905 led to the theory later named Special Relativity (Michelson was awarded the Nobel prize in 1907).

By reaching its goal Microscope will improve the current best tests of UFF-WEP by 2 orders of magnitude, to 10^{-15} . Should the possibility of a non zero violation signal emerge from Microscope data, it will call for urgent checking and more precise space experiments such as GG (which aims at 10^{-17}) might become of interest to space agencies. It is therefore important and timely to firmly establish the role of general relativistic effects

^a Paper to appear on Physical Review D

in high precision satellite tests of UFF-WEP.

Deviations from Newtonian predictions in the motion of orbiting bodies have been investigated since the birth of General Relativity, in the hope to provide observational evidence for the new theory.

In November 1915 Einstein presented to the Prussian Academy of Sciences in Berlin his results on the “Explanation of the perihelion motion of Mercury from the General Theory of Relativity” [15], and one week later “The field equations of gravitation” [16]. A year later De Sitter published: “Planetary motion and the motion of the moon according to Einstein’s theory” [17].

Einstein’s Mercury paper and De Sitter’s work mentioned above deal with non rotating masses. However, all celestial bodies rotate. According to Newton, the gravitational field of a celestial body does not depend on whether the body does rotate or not. Not so in General Relativity. In 1918 Thirring and Lense [18, 19] calculated the secular effects of the rotation of the central body on the orbits of planets and moons (see the English translation and comments on the Thirring and Lense papers by [20]).

Since then, considerable theoretical work has been carried out to include also the rotation of the secondary body, leading to the so-called MPD (Mathisson, Papapetrou, Dixon) equations [21–23]. The whole subject was revisited in the early 1960s [24–26] with the proposal to fly a high precision gyroscope in low Earth orbit in order to measure general relativistic effects (the GP-B mission, launched in 2004 [27]) and in the 1970s became of primary importance for understanding binary systems made by very compact rapidly rotating stars [28, 29].

Similarly to the modern torsion balances used for testing UFF-WEP, the Microscope and GG satellites are designed to rotate in order to up-convert the target signal to higher frequency where important noise sources are known to be smaller than they are at lower frequencies [30, 31].

We compute the general relativistic effects in these experiments by referring to the literature available in which the spin angular momentum of both the primary and secondary body are taken into account. We refer in particular to the work of Barker and O’Connell [28, 29] (in checking the equation numbers quoted please note that they refer to the primary and secondary body with the numbers ‘2’ and ‘1’ respectively while in this work we do the opposite).

This paper is organized as follows. In Sec. II we estimate the differential accelerations between the test masses as predicted by GR, in absence of spin as well as in the presence of a spinning Earth and of spinning test cylinders. We also recall the Newtonian differential effects due to the quadrupole mass moments of the interacting bodies. Sec. III deals with differential precession of the orbits due to General Relativity (and Newtonian dynamics) and relates them to the differential accelerations which give rise to them and might compete with the violation signal. In Sec. IV the general relativistic

and Newtonian effects on the spin axes of the test cylinders are estimated. In Sec. V we discuss the effects of rotation on the laser interferometry readout of GG. The conclusions, that in all cases general relativistic effects are negligible by and large, are drawn in Sec. VI.

II. GENERAL RELATIVISTIC ACCELERATIONS COMPETING WITH A VIOLATION SIGNAL

Tests of the Universality of Free Fall are quantified by the fractional differential acceleration

$$\eta = \frac{\Delta a}{a} \quad (1)$$

between two test masses of different composition as they fall in the gravitational field of a source body with the average acceleration a (“driving signal”). The physical observable is the differential acceleration Δa of the falling masses relative to each other, pointing to the center of mass of the source body.

For test masses orbiting the Earth inside a low altitude, low eccentricity, sun-synchronous satellite such as Microscope or GG a violation signal ($\Delta a \neq 0$) driven by the Earth would have the orbital frequency while the driving signal a at the denominator is the gravitational acceleration $g(h)$ caused by the Earth at the satellite altitude h ; $h \simeq 630$ km for GG (with $g(h) \simeq 8.1 \text{ ms}^{-2}$) and slightly higher for Microscope.

Up-conversion of the signal to higher frequency (the higher, the better) is regarded by all experimentalists as a crucial asset because thermal and electronic noise are lower at higher frequency (the higher the better). This fact has been demonstrated by the rotating torsion balances, which have been able to reach the thermal noise limit expected at their rotation rate (see [30], Fig. 20).

For this reason both Microscope and GG are designed to rotate, though the way they accomplish it is different due to the different experiment design.

The test masses are (nominally) concentric hollow cylinders in both cases.

A very good coincidence of the centers of mass is crucial because of a major Newtonian effect caused by off-centering. Because of the non uniformity of the gravitational force there is a tidal differential acceleration from the Earth (non zero gravity gradient); it acts at twice the orbital frequency, but there is also a smaller tidal effect, proportional to the eccentricity of the orbit, which acts at the orbital frequency and therefore competes directly with the sought for violation signal. This is a major limiting factor to space tests of UFF, which cannot be totally eliminated because it is impossible to inject the satellite in an exactly circular orbit. Indeed, for high precision tests the centers of mass of the test cylinders must be centered on one another far better than it is typically achieved at the time of launch by construction and mounting.

In Microscope the test cylinders are required to be concentric within $20\ \mu\text{m}$ at launch, and no further adjustment is performed in space. Microscope scientists plan to use the tidal effect at twice the orbit frequency within *a posteriori* data analysis in order to reduce—from the measurement of this effect—the unknown level of off-centering between the test cylinders to within $0.1\ \mu\text{m}$, i.e. a factor 200 better than achieved via hardware at launch (see [9], p. 4). A recent press release by the French space agency reports measured offsets of $25\ \mu\text{m}$ and $33\ \mu\text{m}$ [32] (the two values are likely to refer to the two accelerometers carried by the Microscope satellite, each one with two test cylinders).

In Microscope each cylinder is constrained to move along its symmetry axis (sensitive axis): weak electrostatic coupling along the axis; one degree of freedom; effect of violation signal maximized when the symmetry axis points to the center of mass of the Earth. Any differential effect (including the violation signal, if any) would displace the centers of mass of the two cylinders relative to each other. An active control loop ensures that they remain centered: the control force itself contains the violation signal along with all classical (and GR) differential effects. In order to up-convert the frequency of the signal to higher frequency the sensitive/symmetry axis must rotate relative to the satellite-to-Earth direction, hence rotation must occur around an axis perpendicular to the symmetry axis of the cylinders. However, it is known in classical mechanics that a rotating axisymmetric rigid body is stable to small perturbations only if rotation occurs around the axis whose principal moment of inertia is distinct from the other two. The rotation mode of Microscope is slow and actively controlled, up to a maximum rate of about $\frac{1}{900}$ Hz, with roughly a factor of 7 up-conversion from the orbital/signal frequency in absence of rotation ([9],[33]).

In GG the cylinders are allowed to move in the plane perpendicular to the symmetry axis (sensitive plane), where they respond to any differential acceleration: weak mechanical coupling in the plane; two degrees of freedom; effect of violation signal maximized when the symmetry axis is perpendicular to the orbit plane. The violation signal would displace the test cylinders to a new equilibrium position and the displacement is measured by a readout laser gauge. In accordance with the cylindrical symmetry of the system rotation occurs around the symmetry axis, and it is stable, hence passive attitude stabilization of the satellite is ensured (no active attitude control needed). At the same time the signal is up-converted from the orbital frequency of $1.7 \cdot 10^{-4}$ Hz to the much higher rotation frequency of 1 Hz (with an up-conversion factor of almost 6000) where electronic and thermal noise are much lower [31, 34]. The rotation frequency is provided at the start of the mission and maintained by angular momentum conservation. Hence the whole satellite spins with no need of motor and bearings, which are a well known major source of noise for all rotating experiments in ground laboratories.

A rotating conductor in the magnetic field of the Earth is known to slow down because of energy dissipation due to eddy currents induced in the conductor by a component of the magnetic field perpendicular to the spin axis. In GG the largest such effect will take place in the *Be* test cylinder (the outer shell of the spacecraft will be manufactured in carbon fiber, and the inner *Ti* test cylinder has smaller size and lower conductivity). This effect has been calculated in [34], Sec. IIIB, and found to be extremely small: the *Q* factor of spin energy dissipation is $1.4 \cdot 10^{10}$, which means that in the total 1 year duration of the mission the spin frequency will decrease by about 1%. This value has been obtained in the presence of a magnetic field of the Earth reduced by a factor 150 by means of a μ -metal shield, and under worst case assumptions. Any residual differential rotation between the outer shell of the GG spacecraft and its inner parts will be sensed and compensated, if needed, by means of the cold gas thrusters in charge of compensating for non gravitational forces (mostly drag from residual atmosphere).

Since the cylinders are suspended and coupled very weakly (taking advantage of the absence of weight in orbit), the frequencies of their normal modes are much lower than the spin frequency. This is a dynamical regime known as ‘super-critical rotation’ (spin speed above the normal mode/critical speed), which ensures ‘self-centering’ better than achieved via construction and mounting by as much as the ratio of the spin-to-normal mode frequency squared. It is well known that such self-centering by physical laws requires two degrees of freedom: see Ch. 6 of Den Hartog textbook [35], particularly Eq. (6.2) for self-centering and Fig. 6.4 for evidence of rotation instability for systems with 1 degree of freedom. It cannot therefore be exploited in Microscope.

It is also well known that the presence of non zero internal damping in the rotating system (rotating damping) gives rise to a slowly growing ‘whirling’ motion at the normal mode frequency: the smaller the damping, the weaker the instability, the slower its growth, the smaller the fraction of the suspension force which is required to damp it (see [35] Sec. 7.4; [36] Sec. 4.5, and [37]). In GG self-centering by physical laws is ensured at a few tens of picometers, and whirl is damped by capacitance sensors/actuators so as not to exceed a separation level between the centers of mass of $1.7\ \text{nm}$; a noise well within the reach of capacitance bridges. Whirl damping is off during science data taking so that the test masses are totally passive save for the laser light of the interferometer in charge of reading their differential displacements.

Only differential accelerations between the test masses do compete with the target violation signal. Accelerations caused by the primary body (the Earth) and ascribed to GR have a specific dependence on the orbiting distance of each test mass; its first order differential effect is linear with the offset Δr of the test masses in the direction to the center of mass of the Earth, which may mimic a violation signal.

All effects predicted by GR on the test masses contain the very small dimensionless parameter:

$$\epsilon = \frac{GM}{c^2 r} \simeq 6.3 \cdot 10^{-10} \quad (2)$$

where G is the universal constant of gravity, c the speed of light, M the mass of the Earth and r the orbital distance of the test body around it. The small parameter ϵ is the ratio between the Schwarzschild radius of the Earth $GM/c^2 = 4.4$ mm and the satellite orbital distance $r = R + h \simeq 7 \cdot 10^6$ m.

The largest GR acceleration on each test mass, a_E , was computed by Einstein in 1915 [15] for non rotating interacting bodies (see also [28] Eq. (65a)). The second largest one, a_{S_1} , is due to the spin of the primary body (spin-orbit interaction; see [28] Eq. (65c)) while the smallest one, $a_{S_1 S_2}$, is due to the fact that both the primary body and the test mass are spinning (spin-spin interaction; see [28] Eq. (65d)). Their respective orders of magnitude are:

$$a_E \simeq 4\epsilon g(h) \simeq 2 \cdot 10^{-8} \text{ ms}^{-2} \quad (3)$$

$$a_{S_1} \simeq 6\epsilon \frac{S_1}{M} \frac{v}{r^2} \simeq 5.7 \cdot 10^{-10} \text{ ms}^{-2} \quad (4)$$

($S_1 \simeq 0.33MR^2\omega_1$ is the spin angular momentum of the Earth with angular velocity $\omega_1 \simeq 7.3 \cdot 10^{-5} \text{ rad s}^{-1}$, hence $\frac{S_1}{M} \simeq 9.8 \cdot 10^8 \text{ m}^2 \text{ s}^{-1}$; $v \simeq 7.5 \cdot 10^3 \text{ m s}^{-1}$ is the orbital velocity of the satellite at distance $r \simeq 7 \cdot 10^6$ m) and:

$$a_{S_1 S_2} \simeq 3\epsilon \frac{S_1}{M} \frac{S_2}{m} \frac{1}{r^3} \simeq 4.8 \cdot 10^{-22} \text{ ms}^{-2} \quad (5)$$

with S_2 the spin angular momentum of a test body of mass m . For the hollow test cylinders of GG spinning around the symmetry axis with angular velocity $\omega_2 = 2\pi \text{ rad s}^{-1}$, inner radius a , outer radius b , height H , it is $S_2 = \frac{1}{2}m(a^2 + b^2)\omega_2$ hence, in the worst case (largest value of S_2) of the outer cylinder ($a \simeq 10.5$ cm, $b \simeq 13$ cm, $H \simeq 28.6$ cm), it is $\frac{S_2}{m} \simeq 0.088 \text{ m}^2 \text{ s}^{-1}$.

What matters in UFF/WEP tests is the differential acceleration between the test masses. In the case of the GR effects (3) and (4) a differential acceleration arises because of a non zero offset Δr between the centers of mass of the test cylinders. In the case of GG, with $\Delta r \simeq 1.7$ nm as reported above, we have:

$$\Delta a_E \simeq 3a_E \frac{\Delta r}{r} \simeq 1.5 \cdot 10^{-23} \text{ ms}^{-2} \quad (6)$$

$$\Delta a_{S_1} \simeq \frac{7}{2} a_{S_1} \frac{\Delta r}{r} \simeq 4.8 \cdot 10^{-25} \text{ ms}^{-2} \quad (7)$$

Instead, the acceleration (5) depends on the geometrical properties of the test cylinders, which are necessarily different because they have been designed to be one inside the other, yielding a differential acceleration larger than

the one caused by off-centering. A worst case assumption is for the differential acceleration to be of same order as the the acceleration itself:

$$\Delta a_{S_1 S_2} \simeq a_{S_1 S_2} \simeq 3 \cdot 10^{-22} \text{ ms}^{-2} \quad (8)$$

Should the differential accelerations (6)-(8) not be identified as due to General Relativity, they might be misinterpreted as a violation of UFF/WEP at the corresponding (spurious) levels:

$$\eta_E = \frac{\Delta a_E}{g(h)} \simeq 1.8 \cdot 10^{-24} \quad (9)$$

$$\eta_{S_1} = \frac{\Delta a_{S_1}}{g(h)} \simeq 6 \cdot 10^{-26} \quad (10)$$

$$\eta_{S_1 S_2} = \frac{\Delta a_{S_1 S_2}}{g(h)} \simeq 3.7 \cdot 10^{-23} \quad (11)$$

showing that even the largest one, caused by the spin-spin interaction, is smaller than the GG target $\eta_{GG} = 10^{-17}$ by more than 5 orders of magnitude. There is therefore no need to investigate the specific signature (frequency, phase, dependence on the orbital parameters) and the exact values of these effects which come into play in the framework of General Relativity.

In the case of Microscope the spin-spin effect (11) is even less relevant than it is for GG because of a smaller value of S_2/m (mostly because of a slower spin rate by almost 3 orders of magnitude) and also because of the lower precision target $\eta_{microscope} = 10^{-15}$ of the mission. As for (9) and (10) they are about 60 times larger for Microscope (assuming $\Delta r \simeq 0.1 \mu\text{m}$ as reconstructed a posteriori) hence their ratio to the mission target is about a factor of two smaller than the corresponding one for GG.

It is worth noticing that, as expected, the general relativistic effects considered above are much smaller than the Newtonian ones due to the non-zero quadrupole mass moments of the Earth and the test masses.

The quadrupole mass moment of the Earth $J_2^{(1)} \simeq 10^{-3}$ gives rise to an additional acceleration on each test mass (see e.g. [28], Eq. (65e))

$$a_{Q_1} \simeq \frac{3}{2} g(h) J_2^{(1)} \left(\frac{R}{r} \right)^2 \simeq 10^{-2} \text{ ms}^{-2} \quad (12)$$

If the centers of mass of the test bodies are well centered on one another its differential value is below the target. For GG we have:

$$\Delta a_{Q_1} \simeq 4a_{Q_1} \frac{\Delta r}{r} \simeq 10^{-17} \text{ ms}^{-2} \quad (13)$$

hence

$$\eta_{Q_1} = \frac{\Delta a_{Q_1}}{g(h)} \simeq 1.3 \cdot 10^{-18} \quad (14)$$

which is a factor 8 smaller than the mission target $\eta_{GG} = 10^{-17}$.

If each test cylinder has a non zero quadrupole mass moment because its principal moments of inertia are not all equal, the Earth's monopole does couple with it yielding an additional (Newtonian) acceleration which has been known as a major limitation to tests of UFF/WEP aiming at very high precision [38, 39]. In the specific configuration of the GG experiment, with principal moments of inertia I_z relative to the spin/symmetry axis, $I_x = I_y$ relative to the cartesian axes in the plane perpendicular to it and a non zero value of the ratio $\frac{\Delta I}{I_x} = \frac{I_z - I_x}{I_x}$, this effect has been calculated to be of the order (see [40], Sec. 2.2.5):

$$a_{Q_2} \simeq \frac{3}{8} g(h) \frac{\Delta I}{I_x} \frac{r_Q^2}{r^2} \quad (15)$$

with $r_Q^2 = a^2 + b^2 + \frac{H^2}{3}$. The corresponding differential acceleration is dominated by the different value for the two test cylinders of the factor $\frac{\Delta I}{I_x} r_Q^2$. In GG they have been designed so as to make this effect much smaller than the target signal:

$$\Delta a_{Q_2} \simeq 5.6 \cdot 10^{-18} \text{ ms}^{-2} \quad (16)$$

yielding a (spurious) violation at the level:

$$\eta_{Q_2} = \frac{\Delta a_{Q_2}}{g(h)} \simeq 6.9 \cdot 10^{-19} \quad (17)$$

which is about a factor 14 below the target. Note that this result has been obtained with a fractional difference in the moments of inertia for each test cylinder of the order of 0.01, which is not a demanding requirement at all.

Should GG aim at 10^{-18} with the same level of centering of the test masses, both effects (13) and (16) would be close to the target signal. However, their signature is different from that of the signal and known exactly from celestial mechanics (in the case of (13) the value of J_2 of the Earth is well determined in satellite geodesy). Hence they can be separated from the signal by means of various measurements, each one to the target precision, in different dynamical conditions (e.g. different angles between the spin axis and the normal to the orbit plane). Many such measurements are possible because of the short integration time required by GG [34].

In the case of Microscope a correct estimate of the effect (15) should be calculated taking into account the specific geometry and mass distribution of the test bodies. However, for the target $\eta_{Microscope} = 10^{-15}$ this effect is not a matter of concern, and it has never been listed in the error budget of the mission.

III. GENERAL RELATIVISTIC EFFECTS ON THE ORBITS OF THE TEST MASSES

In a 2-body problem the secular effects due to General Relativity on the semimajor axis of the orbit are zero.

The orbital angular momentum vector (perpendicular to the orbit plane) and the Lenz vector (a vector pointing to the pericenter of the orbit whose modulus is the orbital eccentricity) which within Newtonian gravity are both fixed in inertial space, within GR are subjected to a secular precession with the same angular velocity. Since the precession velocity is the same, the orbit precesses as a whole (see [28], Eqs. (73)-(74)).

All GR contributions to orbit precession are proportional to the ϵ parameter (2). There is a contribution independent of rotation, a contribution due to the rotation of the primary body and a contribution due to the rotation of both the primary and secondary body. We have:

$$\Omega_E^{orbit} \simeq 3\epsilon n \simeq 2 \cdot 10^{-12} \text{ rad s}^{-1} \quad (18)$$

with n the mean orbital angular velocity of the satellite (see [28], Eq. (76a)),

$$\Omega_{S_1}^{orbit} \simeq 0.66\epsilon \left(\frac{R}{r}\right)^2 \omega_1 \simeq 2.5 \cdot 10^{-14} \text{ rad s}^{-1} \quad (19)$$

(see [28], Eq. (76c) with $S_1 \simeq 0.33MR^2\omega_1$) and

$$\begin{aligned} \Omega_{S_1 S_2}^{orbit} &\simeq \frac{3}{4} 0.33\epsilon \left(\frac{R}{r}\right)^2 \frac{a^2 + b^2}{r^2} \frac{\omega_1 \omega_2}{n} \\ &\simeq 3.16 \cdot 10^{-26} \text{ rad s}^{-1} \end{aligned} \quad (20)$$

(see [28], Eq. (76d) with $S_1 \simeq 0.33MR^2\omega_1$ and $S_2 \simeq \frac{1}{2}m(a^2 + b^2)\omega_2$ for the test cylinders of GG and a worst case estimate).

Since UFF/WEP tests are differential experiments, only the differential precession of the orbits of the test cylinders relative to each other is relevant. For the orbit precessions (18) and (19) the difference between the two cylinders is due to the fact that they are not exactly at the same distance from the center of mass of the Earth, hence:

$$\Delta\Omega_E^{orbit} \simeq \frac{5}{2} \Omega_E \frac{\Delta r}{r} \simeq 1.2 \cdot 10^{-27} \text{ rad s}^{-1} \quad (21)$$

and

$$\Delta\Omega_{S_1}^{orbit} \simeq 3\Omega_{S_1}^{orbit} \frac{\Delta r}{r} \simeq 1.8 \cdot 10^{-29} \text{ rad s}^{-1} \quad (22)$$

$$\begin{aligned} \Delta\Omega_{S_1 S_2}^{orbit} &\simeq \frac{3}{4} 0.33\epsilon \left(\frac{R}{r}\right)^2 \frac{\omega_1 \omega_2}{n} \frac{1}{r^2} (\Delta(a^2 + b^2)) \\ &\simeq 2 \cdot 10^{-26} \text{ rad s}^{-1} \end{aligned} \quad (23)$$

In addition, it has been shown ([28], Eq.(46)) that according to GR the quadrupole mass moment of the primary body gives an additional contribution to the precession of the orbit of the test mass:

$$\Omega_{Q_1 GR}^{orbit} \simeq \frac{9}{4} \epsilon J_2^{(1)} \left(\frac{R}{r}\right)^2 n \simeq 1.4 \cdot 10^{-15} \text{ rad s}^{-1} \quad (24)$$

The corresponding differential precession between the orbits of the test masses, with the center of mass offset as in the case of GG, is:

$$\Delta\Omega_{Q_1GR}^{orbit} \simeq \frac{7}{2}\Omega_{Q_1GR}\frac{\Delta r}{r} \simeq 1.17 \cdot 10^{-30} \text{ rad s}^{-1} \quad (25)$$

These results show that the largest differential precession predicted by General Relativity is $\Delta\Omega_{S_1S_2}^{orbit}$, due to the proper rotation of the Earth coupling with the proper rotation of the test cylinders. However, the angular velocity of differential precession is extremely small, with a period more than 8 orders of magnitude longer than the age of the universe.

By comparison with the GR effects computed above it is worth recalling the Newtonian contributions to orbit precession which are already taken into account in the experiments and their numerical simulations. The largest one is the well known precession due to the quadrupole mass moment of the Earth on any point mass moving on an inclined orbit around it. This effect is exploited by both Microscope and GG in order to keep the satellite in a sun-synchronous orbit (with an appropriate choice of the inclination for the selected altitude). It is given by:

$$\Omega_{Q_1}^{orbit} \simeq \frac{3}{2}J_2^{(1)}\left(\frac{R}{r}\right)^2 n \simeq 1.4 \cdot 10^{-6} \text{ rad s}^{-1} \quad (26)$$

yielding a much smaller differential precession between the orbits of the test masses:

$$\Delta\Omega_{Q_1}^{orbit} \simeq \frac{7}{2}\Omega_{Q_1}\frac{\Delta r}{r} \simeq 1.2 \cdot 10^{-21} \text{ rad s}^{-1} \quad (27)$$

It has been shown in [29], Eq. (71) that an additional Newtonian precession of the orbit occurs if the secondary body has a non zero quadrupole mass moment of its own, coupling with the monopole of the Earth. In this case the dependence on the average size of the test body (squared) makes the effect many orders of magnitude smaller than the previous one. With the typical numbers of GG we have, for the largest body:

$$\Omega_{Q_2}^{orbit} \simeq \frac{3}{2}J_2^{(2)}\left(\frac{\bar{r}}{r}\right)^2 n \simeq 6.4 \cdot 10^{-21} \text{ rad s}^{-1} \quad (28)$$

where $J_2^{(2)}$ is the quadrupole mass moment of the test body (defined similarly to the quadrupole mass moment $J_2^{(1)}$ of the Earth in the expansion of its gravitational field in multipole mass moments) and \bar{r} is the average size (half the sum of the inner and outer radius of the test cylinder). The numerical estimate refers to GG (worst case value), and the corresponding differential precession turns out to be only one order of magnitude smaller:

$$\Delta\Omega_{Q_2}^{orbit} \simeq 7.5 \cdot 10^{-22} \text{ rad s}^{-1} \quad (29)$$

a value slightly smaller than the differential precession (27) due to the quadrupole mass moment of the Earth.

As we can see by comparing (18) and (26), the largest precession due to GR is 6 orders of magnitude smaller than the largest Newtonian precession, while in the case of differential precession the Newtonian ones dominate by 4 orders of magnitude (see (27) and (29) in comparison with the largest GR differential precession (23)).

In order to assess how much these orbit precessions affect the test we use the variation of the elements perturbative equations in the form of Gauss (see [41]), in which the time variation of the orbital elements are expressed in terms of the radial, transverse and out-of-plane component of the perturbing acceleration a_R , a_T and a_W respectively. For the effects on the pericenter and the node we use Eqs, (3.43) and (3.47) of [41] and find the following relationships between a (differential) orbit precession rate and the corresponding (differential) components of the perturbing acceleration that generates it:

$$\Delta a_W \simeq v\Delta\Omega_{Q_2}^{orbit} \simeq 5.6 \cdot 10^{-18} \text{ m s}^{-2} \quad (30)$$

$$\Delta a_R \simeq \Delta a_T \simeq ev\Delta\Omega_{Q_2}^{orbit} \simeq 5.6 \cdot 10^{-20} \text{ m s}^{-2} \quad (31)$$

As expected, they are related through the orbital velocity v , and –for the same precession of the orbit– the radial and transverse components of the perturbation involved are smaller than the out-of-plane component by as much as the orbital eccentricity e (we have used the maximum value required for GG $e \lesssim 0.01$; for Microscope the requirement is $e \lesssim 0.005$).

Only the radial component Δa_R would compete with the violation signal. The corresponding spurious contribution to violation comes from the Newtonian differential precession and amounts to:

$$\eta_{orbit-precession} = \frac{\Delta a_R}{g(h)} \simeq \frac{5.6 \cdot 10^{-20}}{8.1} \simeq 6.9 \cdot 10^{-21} \quad (32)$$

which is more than 3 orders of magnitude smaller than the GG target. As for the contribution from the largest GR effect (22), it is 4 orders of magnitude smaller still.

IV. GENERAL RELATIVISTIC EFFECTS ON THE SPIN AXES OF THE TEST MASSES

In the framework of General Relativity the spin axes of the test masses are subjected to the following precessions:

$$\Omega_E^{spinaxis} \simeq \frac{3}{2}\epsilon n \simeq 10^{-12} \text{ rad s}^{-1} \quad (33)$$

caused by the primary body (the Earth) regardless of its rotation (also known as De Sitter precession; see e.g. [28], Eq. (42)), and

$$\Omega_{S_1}^{spinaxis} \simeq \frac{0.33}{2}\epsilon\left(\frac{R}{r}\right)^2 \omega_1 \simeq 6.3 \cdot 10^{-15} \text{ rad s}^{-1} \quad (34)$$

due to the proper rotation of the Earth with spin angular momentum $S_1 \simeq 0.33MR^2\omega_1$ (also known as Lense-Thirring precession; see e.g [28] Eq. (29)). The corresponding differential precessions between the spin axes of the two cylinders, due to the fact that they are not exactly centered on each other, are (GG case):

$$\Delta\Omega_E^{spinaxis} \simeq \frac{5}{2}\Omega_E^{spinaxis}\frac{\Delta r}{r} \simeq 6.2 \cdot 10^{-28} \text{ rad s}^{-1} \quad (35)$$

and

$$\Delta\Omega_{S_1}^{spinaxis} \simeq 3\Omega_{S_1}^{spinaxis}\frac{\Delta r}{r} \simeq 4.6 \cdot 10^{-30} \text{ rad s}^{-1} . \quad (36)$$

A by far larger precession of the spin axes of the test cylinders is Newtonian, due to the fact that they have a non zero fractional difference of their principal moments of inertia $\frac{I_z - I_x}{I_x} = \frac{\Delta I}{I_x}$, z being the direction of the symmetry/rotation axis. According to [29], Eq. (47), the precession rate of the spin axis is:

$$\Omega_{Q_2S_2}^{spinaxis} \simeq \frac{1}{2}\frac{\Delta I}{I_x}\frac{n^2}{\omega_2} \simeq 1.3 \cdot 10^{-9} \text{ rad s}^{-1} \quad (37)$$

where the numerical estimate refers to the GG test cylinders (worst case: $\frac{\Delta I}{I_x} \simeq 0.014$). Note that this angular precession rate is proportional to the ratio $\frac{n^2}{\omega_2}$ between the orbital mean motion n squared and the rotation angular velocity of the test cylinder ω_2 , thus implying that for test cylinders with the same orbital velocity those which spin faster have a slower precession rate (the ratio of the spin rates is about 900 to 1 between GG and Microscope). The corresponding differential precession between the test cylinders is not much smaller because they cannot have values of $\frac{\Delta I}{I_x}$ exactly (or very nearly) equal. In GG, by requiring that the difference is of $2 \cdot 10^{-3}$ the differential precession rate of the spin axes is:

$$\Delta\Omega_{Q_2S_2}^{spinaxis} \simeq \frac{1}{2}\Delta\left(\frac{\Delta I}{I_x}\right)\frac{n^2}{\omega_2} \simeq 1.8 \cdot 10^{-10} \text{ rad s}^{-1} \quad (38)$$

with a differential precession period of one thousand years. In the planned one year duration of the mission the differential precession angle amounts to about 0.3° while each spin axis precesses by about 2.4° . We recall that in GG the test cylinders spin around their symmetry axes, hence the rotation is stable against small perturbations. In addition, the spin frequency is higher than the normal mode frequencies in the plane perpendicular to the spin/symmetry axis: a dynamical condition which is known to ensure natural damping of the conical modes (precessions) (see [36], [42]).

A realistic estimate of spin axes precessions in the case of Microscope would require knowledge of the geometrical and mass properties of the test cylinders and of the rotation control of the system. However, it is apparent that the Newtonian framework under which the experiment has been designed is fully adequate to the task, since precessions due to General Relativity are many many orders of magnitude smaller than the Newtonian ones.

V. EFFECTS ON THE READOUT LASER GAUGE

In GG the relative displacements of the test cylinders in the sensitive plane perpendicular to the spin/symmetry axis are read by a laser interferometry gauge. It has been proposed in 2010 by M. Shao (JPL) in substitution of the originally planned capacitance readout (tested in the laboratory prototype of GG) because of its numerous advantages [43]. A violation signal at the target level of 10^{-17} shows up as a 0.6 pm displacement between the centers of mass of the test cylinders pointing to (or away from) the center of mass of the Earth at the spin frequency of 1 Hz (after up-conversion by rotation from the much lower orbital frequency of $1.7 \cdot 10^{-4}$ Hz). The laser gauge is expected to have a displacement noise of $\frac{1 \text{ pm}}{\sqrt{\text{Hz}}}$ at 1 Hz.

In a spinning experiment the readout is obviously co-rotating with the system. As reported in Sec. I, concerns have been expressed by the SARP panel of ESA, about the effects of rotation (of the tests cylinders and the Earth) on the readout laser gauge.

The laser gauge designed for GG is presented in [44] where some key sources of noise are discussed on the basis of the results of specific lab tests. In [45] a measured displacement noise of $\frac{3 \text{ pm}}{\sqrt{\text{Hz}}}$ is reported at 1 Hz and the onset of a spurious displacement in the presence of rotation (Sagnac effect) is discussed and quantified. The issue is as follows: if –from the point of separation to the point of recombination– the interfering laser rays happen to enclose a non-zero area, and the axis normal to this area has a component along the the rotation axis, then the laser rays traveling in the sense of rotation and those traveling opposite to it (both at the speed of light) do have different flight times, yielding a spurious interference signal. In laser gyros this ‘spurious’ signal is used to measure the rotation angular velocity perpendicular to the area enclosed by the gyro (e.g. the diurnal rotation velocity the Earth).

In the case of GG the laser rays are (nominally) aligned, and the area enclosed from separation to recombination is zero, hence there should be no spurious displacement. However, a non-zero area arises in the presence of a misalignment. As shown in [45], a misalignment of $10 \mu\text{m}$ (with a typical 20 cm separation from beam launcher to target) results in a spurious displacement of $\simeq 4 \cdot 10^{-14}$ m. It is 2 orders of magnitude smaller than the signal; moreover, only its time variation at the spin frequency does compete with the violation signal (a constant bias doesn’t matter). There is even a better way out: the lasers can be arranged in such a way that the angular velocity vector of rotation has no component perpendicular to the non-zero area resulting from the misalignment and therefore there is (nominally) no Sagnac effect. In reality, the effect is reduced even further. This strategy has been followed from the start in the design of the GG laser gauge.

As far as the effects of the rotation of the Earth on

the laser gauge of GG are concerned, it is known that the rotation of a celestial body (and also its flattening) do affect the angle of deviation and the propagation time of light rays which, in their journey from emitter to receiver happen to pass close to the body (see e.g. [46]). In high precision astronomical measurements such as those carried out by the GAIA mission of ESA these effect are indeed carefully calculated and taken into account in the framework of General Relativity. However, in GG there is no measurement over astronomical distances: the laser interferometer works inside the spacecraft, with optical path differences (in between the targets located, respectively, on the outer surface of the inner cylinder and on the inner surface of the outer one) of 2 cm or less, and a path length of about 20 cm.

We therefore don't expect that any effect from the spin (and flattening) of the Earth on the laser interferometry readout of GG should be taken into account. More importantly, there is recent experimental evidence which supports this conclusion. A heterodyne laser gauge similar to the one of GG, though more demanding and complex, has recently flown on LISA pathfinder (LPF) [47]. In this case the optical path difference is $\simeq 38$ cm, which is the separation distance between the test masses that constitute the mirrors which, in the final gravitational wave interferometer LISA, will be located in different spacecraft about 5 million km away from each other. The laser gauge of LPF has been designed to achieve low noise down to $7 \cdot 10^{-4}$ Hz (while in GG the violation signal is up-converted by rotation to 1 Hz). Reaching this level of noise at such low frequencies and over 38 cm separation of the test masses requires –among other things– the frequency of the laser to be stabilized. This is a demanding requirement in space, which does not apply in the case of GG because of the 1 Hz frequency of the signal and 2 cm maximum separation. The displacement noise as measured by LPF above 0.06 Hz is reported to be of $\frac{0.035 \text{ pm}}{\sqrt{\text{Hz}}}$ (about 30 times lower than required for GG at 1 Hz); it

is interpreted by the authors as entirely due to the interferometer and no evidence is reported of any effect due to the rotation and flattening of the Earth [47]. In addition, it turns out to be about 100 times lower in absence of weight than obtained in ground tests before launch.

VI. CONCLUSIONS

The Microscope satellite, launched on 25 April 2016, is testing the Universality of Free Fall and the Weak Equivalence Principle in the gravitational field of the Earth aiming at two orders of magnitude improvement over the current best tests. In a similar orbit but with a different experiment design, the GG satellite aims at a test one hundred times better than Microscope. They must detect extremely small differential accelerations acting between test cylinders of different composition, ruling out any competing effect which is due to known physics.

In 2015, during the evaluation process of the space mission proposals shortlisted as candidates for the medium size mission M4 of the European Space Agency, the panel appointed by the Agency to evaluate GG made the point that while the experiment has been designed in the framework of Newtonian physics, it should instead take into account general relativistic effects, in particular those which involve the spin angular momentum of the source body (the Earth) and that of the test cylinders.

We have carefully analyzed all known general relativistic effects on the test cylinders of GG and Microscope showing that they are all negligible by and large.

Acknowledgements. Thanks are due to Norbert Wex, David Lucchesi, Angelo Tartaglia, Slava Turyshev, Juergen Mueller, Alberto Anselmi and Maria Teresa Crosta for their contributions. Support from ESA in the development of a low noise laser gauge over small distances is gratefully acknowledged.

-
- [1] A. M. Nobili, D. M. Lucchesi, M. T. Crosta, M. Shao, S. G. Turyshev, R. Peron, G. Catastini, A. Anselmi, and G. Zavattini, On the universality of free fall, the equivalence principle, and the gravitational redshift, *Am. J. Phys.* 81, 527 (2013)
 - [2] A. Einstein, *Grundlage der allgemeinen Relativitätstheorie*, *Annalen der Physik* (ser. 4), 49, 769-822 (1916); *The foundation of the general theory of relativity*, in *The Principle of Relativity*, Dover Publ. Inc. N.Y., USA, 1952)
 - [3] S. Baeßler, B. R. Heckel, E. G. Adelberger, J. H. Gundlach, U. Schmidt, and H. E. Swanson, Improved Test of the Equivalence Principle for Gravitational Self-Energy, *Phys. Rev. Lett.* 83, 3585 (1999)
 - [4] S. Schlamminger, K.-Y. Choi, T. A. Wagner, J. H. Gundlach, and E. G. Adelberger, Test of the Equivalence Principle Using a Rotating Torsion Balance, *Phys. Rev. Lett.* 100, 041101 (2008)
 - [5] T. A. Wagner, S. Schlamminger, J. Gundlach, and E. G. Adelberger, Torsion-balance tests of the weak equivalence principle, *Class. Quantum Grav.* 29, 184002 (2012)
 - [6] J. G. Williams, S. G. Turyshev & D. H. Boggs, Progress in Lunar Laser Ranging Tests of Relativistic Gravity, *Phys. Rev. Lett.* 93, 261101 (2004)
 - [7] J. G. Williams, S. G. Turyshev & D. H. Boggs, Lunar laser ranging tests of the equivalence principle, *Class. Quantum Grav.* 29, 184004 (2012)
 - [8] J. Mueller, F. Hoffman & L. Biskupek, Testing various facets of the equivalence principle using lunar laser ranging, *Class. Quantum Grav.* 29, 184006 (2012)
 - [9] P. Touboul, G. Metris, V. Lebat, and A. Robert, The MICROSCOPE experiment, ready for the in-orbit test of the equivalence principle, *Class. Quantum Grav.* 29, 184010 (2012)
 - [10] A. M. Nobili, M. Shao, R. Pegna, G. Zavattini, S. G. Turyshev, D. M. Lucchesi, A. De Michele, S. Doravari, G.

- L. Comandi, T. R. Saravanan, F. Palmonari, G. Catastini and A. Anselmi, ‘Galileo Galilei’ (GG): space test of the weak equivalence principle to 10^{-17} and laboratory demonstrations, *Class. Quantum Grav.* 29, 184011 (2012)
- [11] Lin Thou, Shitong Long, Biao Tang, Xi Chen, Fen Gao, Wencui Peng, Witao Duan, Jiaqi Zhong, Zongyuan Xiong, Jin Wang, Yaunzhong Zhang, and Mingsheng Zhan, Test of Equivalence Principle at 10^{-8} Level by a Dual-Species Double-Diffraction Raman Atom Interferometer, *Phys. Rev. Lett.* 115, 013004 (2015)
- [12] A. M. Nobili, Fundamental limitations to high-precision tests of the universality of free fall by dropping atoms, *Phys. Rev. A.* 93, 023617 (2016)
- [13] G. Arnold, A. Bazzano, P. Binetruy, E. Chassefiere, M. Diaz Trigo, J. Drew, E. Friis-Christensen, O. Grasset, G. Harendel, C. Haswell, M. Hewitson, G. Nelemans, B. Nordstrom, W. Percival, G. Peres, J. Rodriguez-Pacheco Martin, M. Sauvage, B. Stelzer, E. Szuszkiewicz, R. Tucker, J. Zarnecki, Report to ESA on the GG proposal for the medium size mission M4 (2015)
- [14] A. A. Michelson and E. W. Morley, On the relative motion of the Earth and the luminiferous ether, *Am. J. Science* 34, 333 (1887)
- [15] A. Einstein, Erklärung der Perihelwegung Merkur der allgemeinen Relativitätstheorie Königlich Preussische Akademie der Wissenschaften, Berlin, p. 831-839 (1915). See also: Explanation of the perihelion motion of Mercury from the General Theory of Relativity, in *The collected papers of Albert Einstein. Volume 6*, A. J. Kox, M. J. Klein, and R. Schumann, Editors. Princeton University Press (1996).
- [16] A. Einstein, Die Feldgleichungen der Gravitation, Preussische Akademie der Wissenschaften, Sitzungsberichte part 2, 844-847 (1915); The field equations of gravitation, http://en.wikisource.org/wiki/The_Field_Equations_of_Gravitation
- [17] W. De Sitter, Planetary motion and the motion of the moon according to Einstein’s theory, *Amsterdam Proc.* 19 (1916)
- [18] H. Thirring, Über die Wirkung rotierender ferner Massen in der Einsteinschen Gravitationstheorie (On the effect of rotating distant masses in Einstein’s theory of gravitation), *Phys. Z.* 19, 33 (1918)
- [19] J. Lense and H. Thirring, Über den Einfluss der Eigenrotation der Zentralkörper auf die Bewegung der Planeten und Monde nach der Einsteinschen Gravitationstheorie (On the influence of the proper rotation of central bodies on the motions of planets and moons according to Einstein’s theory of gravitation), *Phys. Z.* 19, 156 (1918)
- [20] B. Mashhoon, F. W. Hehl, and D. S. Theiss, On the Gravitational Effects of Rotating Masses: The Thirring-Lense Papers, *Gen. Relat. Grav.* 16, 711-750 (1984)
- [21] M. Mathisson, Neue Mechanik materieller Systeme, *Acta Phys. Pol.* 6, 163 (1937); rpublished as *New mechanics of material systems*, *Gen. Relativ. Gravit.* 42, 1011 (2010)
- [22] A. Papapetrou, Spinning test-particles in General Relativity I. *Proc. Roy. Soc. Lond.* A209, 248 (1951)
- [23] W. G. Dixon, A covariant multipole formalism for extended test bodies in General Relativity, *Nuovo Cimento* 34, 317 (1964)
- [24] L. Schiff, Possible new experimental tests of general relativity theory, *Phys. Rev. Lett.* 4, 215-217 (1960)
- [25] L. Schiff, Motion of a gyroscope according to Einstein’s theory of gravitation, *Proc. Nat. Acad. Sci.*, 46, 871-882 (1960)
- [26] L. Schiff, Proceedings of the Conference on “Experimental tests of theories of relativity”, Stanford (1961)
- [27] C. W. F. Everitt, D. B. Debra, B. W. Parkinson, J. P. Turneaure, J. W. Conklin, M. I. Heifetz, G. M. Keiser, A. S. Silbergleit, T. Holmes, J. Kolodziejczak, M. Al-Meshari, J. C. Mester, B. Muhlfelder, V. G. Solomonik, K. Stahl, and P. W. Worden, Jr., W. Bencze, S. Buchman, B. Clarke, A. Al-Jadaan, H. Al-Jibreen, J. Li, J. A. Lipa, J. M. Lockhart, B. Al-Suwaidan, M. Taner, and S. Wang, Gravity Probe B: Final Results of a Space Experiment to Test General Relativity, *Phys. Rev. Lett.* 106, 221101 (2011)
- [28] B. M. Barker & R. F. O’Connell, Derivation of the Equations of Motion of a Gyroscope from the Quantum Theory of Gravitation, *Phys. Rev. D* 2, 1428-1435 (1970)
- [29] B. M. Barker & R. F. O’Connell, Gravitational two-body problem with arbitrary masses, spins and quadrupole moments, *Phys. Rev. D* 12, 329-335 (1975)
- [30] E. G. Adelberger, J. H. Gundlach, B. R. Heckel, S. Hoedl, and S. Schlamminger, Torsion balance experiments: A low-energy frontier of particle physics, *Prog. Part. Nucl. Phys.* 62, 102 (2009)
- [31] R. Pegna, A. M. Nobili, M. Shao, S. G. Turyshev, G. Catastini, A. Anselmi, R. Spero, S. Doravari, G. L. Comandi and A. De Michele, Abatement of Thermal Noise due to Internal Damping in 2D Oscillators with Rapidly Rotating Test Masses, *Phys. Rev. Lett.* 107, 200801 (2011)
- [32] CNES-ONERA Press Release, 29 September (2016)
- [33] P. Touboul, Talk given at Microscope Colloquium IV, Paris (2015) <http://microscope4.sciencesconf.org/81163>
- [34] A.M. Nobili, R. Pegna, M. Shao, S. G. Turyshev, G. Catastini, A. Anselmi, R. Spero, S. Doravari, G. L. Comandi, D. M. Lucchesi, and A. De Michele, Integration time in space experiments to test the equivalence principle, *Phys. Rev. D* 89, 042005 (2014)
- [35] J. P. Den Hartog, *Mechanical vibrations* (Dover Publications Inc., New York 1985; first published 1934).
- [36] G. Genta, *Vibrations of structures and machines* (Springer-Verlag, New York 1993)
- [37] A. M. Nobili, D. Bramanti, E. Polacco, G. Catastini, A. Anselmi, S. Portigliotti, A. Lenti, P. di Giamberardino, S. Monaco, and R. Ronchini, Evaluation of a proposed test of the weak equivalence principle using Earth-orbiting bodies in high-speed co-rotation: re-establishing the physical bases, *Class. Quantum Grav.* 16, 1463-1470 (1999)
- [38] J. P. Blaser *et al.*, *Satellite Test of the Equivalence Principle*, Report on the Phase A Study, ESA/NASA SCI (93)4 (1993)
- [39] J. P. Blaser *et al.*, *Satellite Test of the Equivalence Principle*, Report on the Phase A Study, ESA SCI (96)5 (1996)
- [40] *GG Phase A Study*, ASI (1998, revised 2000) <http://eotvos.dm.unipi.it/ggweb/phaseA>
- [41] A. Milani, A. M. Nobili, and P. Farinella, *Non-gravitational perturbations and satellite geodesy* (Adam Hilger Ltd., Bristol 1987)
- [42] A. M. Nobili, D. Bramanti, G. Catastini, E. Polacco, G. Genta, E. Brusa, V. P. Mitrofanov, A. Bernard, P. Touboul, A. J. Cook, J. Hough, I. W. Roxburgh, A.

- Polnarev, W. Flury, F. Barlier, C. Marchal, Proposed noncryogenic, nondrag-free test of the equivalence principle in space, *New Astronomy* 3, 175 (1998)
- [43] M. Shao, SIM-like laser gauge and adaptation to GG, talk given at GG Workshop, Pisa 10-12 February 2010
- [44] M. Pisani, G. Mana, and A. M. Nobili, Design of an interferometric displacement sensor with picometer resolution for the Galileo Galilei mission, IEEE Conference Publications, Metrology for Aerospace (MetroAeroSpace), IEEE 978-1-4799-7569-3/15, pp. 591-595 DOI: 10.1109/MetroAeroSpace.2015.7180724 (2015)
- [45] M. Pisani, M. Zucco, and A. M. Nobili, Laboratory tests of a high-precision laser interferometry readout for the GG experiment in space, IEEE Conference Publications, Metrology for Aerospace (MetroAeroSpace), *in press* (2016)
- [46] S. A. Klioner, Influence of the quadrupole field and rotation of objects on light propagation, *Astron. Zh.* 1046 (1991)
- [47] M. Armano *et al.* (119 authors), Sub-Femto-g Free Fall for Space-Based Gravitational Wave Observatories: LISA Pathfinder Results, *Phys. Rev. Lett.* 116, 231101 (2016)